
What would you do with more GOOD ON time?

Ask about the only once-nightly treatment that gives you more GOOD ON time in your day by reducing both OFF time and dyskinesia.



WHAT IS GOCOVRI?

GOCOVRI® (amantadine) extended release capsules is a prescription medicine used:

- for the treatment of dyskinesia (sudden uncontrolled movements) in people with Parkinson's disease who are treated with levodopa therapy or levodopa therapy with other medicines that increase the effects of dopamine in the brain.
- with levodopa and carbidopa in people with Parkinson's disease who are having "off" episodes.

It is not known if GOCOVRI is safe and effective in children.

IMPORTANT SAFFTY INFORMATION

DO NOT take GOCOVRI if you have severe kidney problems.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information (in pocket) or visit Gocovri.com.



As Parkinson's disease (PD) progresses, motor complications can disrupt your daily life.

Within 5 years of diagnosis, up to 50% of people taking levodopa experience OFF episodes, dyskinesia, or both.



OFF time is when your medication isn't working as it should to manage your symptoms.

COMMON OFF SYMPTOMS









Tremor

Muscle stiffness

Slowness

Walking or balance problems



Dyskinesia is side effect of taking Parkinson's medication to control OFF time. You may develop involuntary, erratic, and irregular movements, which can impact you physically, socially, and emotionally.

DYSKINESIA SYMPTOMS









Body swaying

Dance-like movements

Head-bobbing

Writhing or twisting


A goal of treatment is to increase the amount of GOOD ON time each day



GOOD ON time is when you are experiencing symptom control. You are not OFF and not bothered by dyskinesia. You can go about your day with minimal motor symptoms and get back to things you enjoy, with fewer interruptions.

Over time, even as your doctor adjusts your medicine, you may experience more OFF and/or more dyskinesia. These "motor complications" in your day impact the amount of GOOD ON time you're able to have.

REAL PERSPECTIVES



Learn how Kendall found a treatment to help with both his dyskinesia and OFF time



2

For more GOOD ON time, ask about GOCOVRI

GOCOVRI is the only once-nightly medication proven to reduce both dyskinesia and OFF time for increased GOOD ON time in your day.

STUDY RESULTS:

Participants taking GOCOVRI reported improvements from the beginning to the end of the studies.

Primary endpoint



41%
DECREASE IN DYSKINESIA

vs. 14% with placebo

Secondary endpoint



21% DECREASE IN OFF TIME

vs. 15% increase with placebo

Secondary endpoint



3.8-HOUR INCREASE IN GOOD ON TIME

vs. 1.4 hours with placebo

GOCOVRI was studied in 2 clinical trials. Study participants were adults with Parkinson's disease taking levodopa, and many were also taking other PD medications. Participants used diaries to track their symptoms throughout the day.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information (in pocket) or visit Gocovri.com.





What would you do with more GOOD ON time in your day?

IMPORTANT SAFETY INFORMATION (cont'd) WHAT SHOULD I AVOID WHILE TAKING GOCOVRI?

Do not stop or change the dose of GOCOVRI before talking with your doctor. Call your healthcare provider if you have symptoms of withdrawal such as fever, confusion, or severe muscle stiffness.

Do not drink alcohol while taking GOCOVRI as it can increase your chances of serious side effects.

Do not drive, operate machinery, or do other dangerous activities until you know how GOCOVRI affects you.

4

Could GOCOVRI be right for you?

You don't have to compromise GOOD ON time by living with dyskinesia or OFF time. Talk to your healthcare provider about GOCOVRI, the only medication proven to reduce both with 1 nightly dose.

Make the most of your time with your healthcare provider



Track your daily symptoms in a diary



Record your symptoms on video

Check the boxes that apply to start a conversation with your healthcare provider



- ☐ How can I tell what is dyskinesia versus tremor?
- ☐ It's hard to keep a regular schedule because I don't know when I will be OFF, ON, or with dyskinesia.
- Dyskinesia and/or OFF is interfering with my daily activities.



Find out if GOCOVRI is right for you Scan the QR code or visit GOCOVRI.com.

IMPORTANT SAFETY INFORMATION (cont'd)

If you took too much GOCOVRI, call your doctor or go to the nearest hospital emergency room right away.

WHAT ARE THE POSSIBLE SIDE EFFECTS OF GOCOVRI?

- Falling asleep during normal activities. Activities may include driving, talking, or eating. You may fall asleep without being drowsy or warning.
- Suicidal thoughts or actions and depression. Tell your doctor if you have new or sudden changes in mood, behaviors, thoughts, or feelings, including thoughts about hurting yourself or ending your life.
- **Hallucinations.** GOCOVRI can cause or worsen hallucinations (seeing or hearing things that are not real) or psychotic behavior.
- Feeling dizzy, faint or lightheaded, especially when you stand up (orthostatic hypotension). Lightheadedness or fainting may happen when getting up too quickly after long periods of time, when first starting GOCOVRI, or if your dose has been increased.
- **Unusual urges.** Examples include gambling, sexual urges, spending money, binge eating, and the inability to control them.

The most common side effects of GOCOVRI include dry mouth, swelling of legs and feet, constipation, and falls. If you or your family notices that you are developing any new, unusual or sudden changes in behavior or related symptoms, tell your healthcare provider right away.

These are not all the possible side effects of GOCOVRI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

TELL YOUR DOCTOR ABOUT ALL MEDICAL CONDITIONS, INCLUDING IF YOU:

- have kidney problems.
- have unexpected or unpredictable sleepiness, sleep disorders, or currently take medication to help you sleep or make you drowsy.
- are pregnant or plan to become pregnant or are breastfeeding or plan to breastfeed. GOCOVRI may harm your unborn baby and can pass into your breastmilk.

Tell your doctor about all the medicines you take. Include prescription and overthe-counter medicines, vitamins, and herbal supplements.

Especially tell your doctor if you take medicines like sodium bicarbonate, or have had or are planning to have a live flu vaccination (nasal spray). You can receive the flu vaccination shot but should not get a live flu vaccine while taking GOCOVRI.

Please see accompanying full Prescribing Information and Patient Information (in pocket) or visit Gocovri.com.



R

It's GO time

Ask your doctor if once-nightly GOCOVRI could be right for you.

WHAT IS GOCOVRI?

GOCOVRI® (amantadine) extended release capsules is a prescription medicine used:

- for the treatment of dyskinesia (sudden uncontrolled movements) in people with Parkinson's disease who are treated with levodopa therapy or levodopa therapy with other medicines that increase the effects of dopamine in the brain.
- with levodopa and carbidopa in people with Parkinson's disease who are having "off" episodes.

It is not known if GOCOVRI is safe and effective in children.

IMPORTANT SAFETY INFORMATION

DO NOT take GOCOVRI if you have severe kidney problems.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information and Patient Information (in pocket) or visit Gocovri.com.

GOCOVRI is a registered trademark of Supernus Pharmaceuticals, Inc. All other trademarks are property of their respective owners. © 2023 Supernus Pharmaceuticals, Inc. All rights reserved. GOC.2023-0079.



